

30 YEAR HISTORY OF PUBLIC HEALTH FAILURE

Global Burden of Disease Study 1990-2019. Funded by the Bill & Melinda Gates Foundation

Risk factor Low fruit intake Low vegetable intake Low vegetable intake Low program intake Low program intake Low mut & seed intake Low gram intake Low gram intake Low gram intake Low calcium intake Low calcium intake Low seafood omega-3 intake Low polyunsaturated fat intake High transfat intake High sodium intake METABOLIC RISKS Risk factor High sody-mass index Low book miseral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor NON-DIETARY RELATED RISKS Risk factor Low book miseral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Low pollution (particulate matter) Qzone pollution Logariculate matter) Low program logariculate matter) Low program logariculate matter) Low pollution Logaricula	DIETARY-RELATED RISKS	disease risk rating			risk change	
Low fruit intake Low vegetable intake Low vegetable intake Low whole grain intake Low whole grain intake Low milk intake High red meat intake High sugars-weetened beverages Low fibre intake Low salcium intake Low salcium intake Low salcium intake Low polyunsaturated fat intake High sugars-weetened beverages Low fibre intake Low polyunsaturated fat intake High sugars-weetened beverages Low polyunsaturated fat intake Low polyunsaturated fat intake Low polyunsaturated fat intake High sugars-weetened beverages Low fibre intake Low sugars-weetened to a sugars-weetened beverages Low fibre intake Low sugars-weetened beverages Low fibre intake Low sugars-weetened Low fibre inta		1990	2010	2019	1990-2019	2010-2019
Low vegetable intake Low legume intake Low place intake Low nut & seed intake Low milk intake High red meat intake High processed meat intake High processed meat intake Low golyunsaturated beverages Low fibre intake Low polyunsaturated fat intake High sodium intake Low polyunsaturated fat intake High sodium intake METABOLIC RISKS Risk factor High fasting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction MON-DIETARY RELATED RISKS Risk factor 1990 2010 2019 1990-2019 2019 1990-2019 2019 2019 2019 2019 2019 2019 2019						
Low legume intake Low whole grain intake Low whole grain intake Low milk intake High red meat intake High processed meat intake High sugar-sweetened beverages Low fibre intake Low seafood omega-3 intake Low polyunsaturated fat intake High sodium intake High sodium intake High sodium intake High sodium intake WETABOLIC RISKS Risk factor High fasting plasma glucose High stanisp also glucose High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS RISK factor Unsafe water, sanitation and handwashing Ari pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Ozone pysical activity					000	
Low whole grain intake Low mult a seed intake Low mult intake High red meat intake High processed meat intake High sugar-sweetened beverages Low fibre intake Low acalicum intake Low acalicum intake Low acalicum intake Low acalicum intake High sugar-sweetened beverages Low fibre intake Low polyunsaturated fat intake High sugar-sweetened beverages Low fibre intake Low acalicum intake METABOLIC RISKS Risk factor Ingo 2010 Ingo 2019 In					000	
Low nut & seed intake Low milk intake High red meat intake High processed meat intake High sugar-sweetened beverages Low fibre intake Low calcium intake Low seafood omega-3 intake Low polyunsaturated fat intake High transfat intake High sodium intake WETABOLIC RISKS Risk factor High sodium intake WETABOLIC RISKS Risk factor High sugar sweetened beverages WETABOLIC RISKS Risk factor High sugar sweetened fat intake High attransfat intake High sodium intake WETABOLIC RISKS Risk factor High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction WON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution Lead exposure Intimate partner violence Low physical activity PETER Standard Research and a standard Research and Research					000	0
Low milk intake High red meat intake High red meat intake High sugar-sweetened beverages Low fibre intake Low seafood omega-3 intake Low polyunsaturated fat intake High sodium intake Low polyunsaturated fat intake High transfat intake High stransfat intake High sodium intake METABOLIC RISKS Risk factor High asting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Ari pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Provided the measurage of the second state of						
High red meat intake High processed meat intake High sugar-sweetened beverages Low fibre intake Low calcium intake Low seafood omega-3 intake Low seafood omega-3 intake High transfat intake High sudium intake High sodium intake WETABOLIC RISKS Risk factor High fasting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity PEX. Manual as transmidted as balance of specific partners of spec						
High processed meat intake High sugar-sweetened beverages Low fibre intake Low calcium intake Low seafood omega-3 intake Low polyunsaturated fat intake High transfat intake High sodium intake WETABOLIC RISKS Risk factor High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Ozone pollution Lead exposure Intimate partner violence Low physical activity Ozone positive change One of the processed meat intake O						
High sugar-sweetened beverages Low fibre intake Low calcium intake Low calcium intake Low polyunsaturated fat intake High transfat intake High sodium intake WETABOLIC RISKS Risk factor High fasting plasma glucose High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity VEX. Salest states and st						
Low calcium intake Low seafood omega-3 intake Low polyunsaturated fat intake High transfat intake High sodium intake Olisease risk rating Tisk chanter of the control						⊗ ⊗
Low calcium intake Low seafood omega-3 intake Low polyunsaturated fat intake High transfat intake High sodium intake WETABOLIC RISKS Risk factor High fasting plasma glucose High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor 1990 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 1990-2019 2010 2010 2010						
Low seafood omega-3 intake Low polyunsaturated fat intake High transfat intake High sodium intake WETABOLIC RISKS WETABOLIC RISKS Wisk factor High stating plasma glucose High Loh cloelseterol High systolic blood pressure High systolic blood pressure High systolic blood pressure WELATED RISKS NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure WEY Stewards Statemardina pick of this bid					000	000
Low polyunsaturated fat intake High transfat intake High sodium intake WETABOLIC RISKS Risk factor High fasting plasma glucose High Loholesterol High systolic blood pressure High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Positive change Opesitive chang			00000		Ø	8
High sodium intake High sodium intake METABOLIC Alisease risk rating Risk factor High fasting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction MON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Person and Statemendiate risk Populative change Person and Statemendiate risk Person and Statem					Ø Ø	• • • • • • • • • • • • • • • • • • • •
METABOLIC disease risk rating risk change of the change of	· · · · · · · · · · · · · · · · · · ·					
METABOLIC RISKS Risk factor High fasting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity REV. Colourisk Colourand Instrumediate risk Colourand I						
RISKS Risk factor High fasting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction RON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity						
Risk factor High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction Consequently Consequen		dis	sease risk ra	nting	risk c	hange
High fasting plasma glucose High LDL cholesterol High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction WON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity WEY Counties Intermediate risk of the big	RISKS					
High systolic blood pressure High body-mass index Low bone mineral density Kidney dysfunction NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity	Risk factor	1990	2010	2019	1990-2019	2010-2019
High body-mass index Low bone mineral density Kidney dysfunction Convert a convert	High fasting plasma glucose				$\otimes \otimes \otimes$	⊗ ⊗ ⊗
High body-mass index Low bone mineral density Kidney dysfunction	High LDL cholesterol	• •	• •	• •	00	•
Kidney dysfunction	High systolic blood pressure	• •	• •	• •	8	⊗ ⊗ ⊗
NON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity disease risk rating risk change	High body-mass index	•			⊗ ⊗ ⊗	⊗ ⊗ ⊗
ACON-DIETARY RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity disease risk rating risk change risk change 1990 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 1990-2019 2010 2019 2010 2019 2019 2019 2019	Low bone mineral density	• •	• •	• •	•	•
RELATED RISKS Risk factor Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Unsafe water, sanitation and handwashing Air pollution (particulate matter) Air pollution Air pollut	Kidney dysfunction	• •	• •	• •	88	8
Risk factor 1990 2010 2019 1990-2019 2010 Unsafe water, sanitation and handwashing ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ●		disease risk rating		risk change		
Unsafe water, sanitation and handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity	RELATED RISKS					
handwashing Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Opositive change Positive change	Risk factor	1990	2010	2019	1990-2019	2010-2019
Air pollution (particulate matter) Ozone pollution Lead exposure Intimate partner violence Low physical activity Opositive change Positive change						
Ozone pollution Lead exposure Intimate partner violence Low physical activity						
Lead exposure Intimate partner violence Low physical activity						× × × × × × × × × × × × × × × × × × ×
Intimate partner violence Low physical activity						
Low physical activity						⊗ ⊗
KEY: Olow rick Olintermediate rick Olintermedi						88
	Low physical activity					
number of dots = extent of c	KEY:	● low risk ● ● ● intermediate risk ● ● ● ● high risk			positive change number of dots = extent of change	